



LASIK Eye Drop Directions

These drops are to help reduce the risk of complication from your eye surgery. Use the chart below to help you remember your drops and separate all drops by at least 3 minutes. You should wear clear goggles while you are sleeping for the first week after surgery.



Vigamox or Zymaxid: Start this drop 1 day before surgery and continue until the 1 week post-op visit.

PredForte: Start this drop after surgery and continue as directed for 2 weeks after surgery.

Optive: Use these preservative free hourly for 1 week after surgery and as needed after week 1.

☹☹☹☹ 1 Day Before Surgery

☹☹☹☹ Surgery Day

☹☹☹☹ 1 Day After Surgery

☹☹☹☹ 2 Days After Surgery

☹☹☹☹ 3 Days After Surgery

☹☹☹☹ 4 Days After Surgery

☹☹☹☹ 5 Days After Surgery

☹☹☹☹ 6 Days After Surgery

☹☹☹☹ 7 Days After Surgery
Discontinue Day 8

Surgery Day: Use Drops Every Waking Hour **After Surgery**

☹☹☹☹ 1 Day After Surgery

☹☹☹☹ 2 Days After Surgery

☹☹☹☹ 3 Days After Surgery

☹☹☹☹ 4 Days After Surgery

☹☹☹☹ 5 Days After Surgery

☹☹☹☹ 6 Days After Surgery

☹☹☹☹ 7 Days After Surgery

☹☹ 8 Days After Surgery

☹☹ 9 Days After Surgery

☹☹ 10 Days After Surgery

☹☹ 11 Days After Surgery

☹☹ 12 Days After Surgery

☹☹ 13 Days After Surgery

☹☹ 14 Days After Surgery
Discontinue Day 15

Questions?

Eye Institute Laser Vision Center
2800 Third Street, Rapid City, SD 57701
605-718-3937 or 877-718-3937
www.lasikrapidcity.com